

BALL CONVERGENCE OF POTRA-PTAK-TYPE METHOD WITH
OPTIMAL FOURTH ORDER OF CONVERGENCE

IOANNIS K. ARGYROS* and SANTHOSH GEORGE†

Abstract. We present a local convergence analysis Potra-Ptak-type method with optimal fourth order of convergence in order to approximate a solution of a nonlinear equation. In earlier studies such as [1], [5]–[28] hypotheses up to the fourth derivative are used. In this paper we use hypotheses up to the first derivative only, so that the applicability of these methods is extended under weaker hypotheses. Moreover the radius of convergence and computable error bounds on the distances involved are also given in this study. Numerical examples are also presented in this study.

MSC 2020. 65D10, 65D99.

Keywords. Potra-Ptak-type method, Newton’s method, order of convergence, local convergence.

1. INTRODUCTION

Let $F : D \subseteq S \rightarrow S$ is a nonlinear function, D is a convex subset of S and S is \mathbb{R} or \mathbb{C} . Consider the problem of approximating a locally unique solution x^* of equation

$$(1) \quad F(x) = 0.$$

Newton-like methods are famous for finding solution of (1), these methods are usually studied based on: semi-local and local convergence [3, 4, 20, 21, 22, 24, 26].

Third order methods such as Euler’s, Halley’s, super Halley’s, Chebyshev’s [1]–[28] require the evaluation of the second derivative F'' at each step, which in general is very expensive. That is why many authors have used higher order multipoint methods [1]–[28]. In this paper, we study the local convergence of fourth order method defined for each $n = 0, 1, 2, \dots$ by

$$\begin{aligned} y_n &= x_n - F'(x_n)^{-1}F(x_n) \\ z_n &= x_n - F'(x_n)^{-1}(F(x_n) + F(y_n)) \end{aligned}$$

*Department of Mathematical Sciences, Cameron University, Lawton, OK 73505, USA, e-mail: iargyros@cameron.edu.

†Department of Mathematical and Computational Sciences, NIT Karnataka, India-575 025, e-mail: sgeorge@nitk.ac.in.

$$(2) \quad x_{n+1} = x_n - F(x_n)^{-2} F'(x_n)^{-1} F(y_n)^2 (2F(x_n) + F(y_n)),$$

where x_0 is an initial point. Method (2) was studied by Cordero et.al. in [13]. In particular the fourth order of convergence was shown under hypotheses reaching up to the fourth derivative of function F . Notice that method (2) involves three functional evaluations. Therefore the efficiency index $EI = p^{\frac{1}{m}}$ where p is the order of convergence and m is the number of functional evaluations per step gives $EI = 4^{\frac{1}{3}} = 1.5874$. Kung and Traub conjecture [28] that the order of convergence of any multipoint method without memory cannot exceed the bound 2^{m-1} (called the optimal order). Thus, the optimal order for a method with three function evaluations per step should be four.

Other single and multi-point methods can be found in [2, 3, 20, 25] and the references therein. The local convergence of the preceding methods has been shown under hypotheses up to the fourth derivative (or even higher). These hypotheses restrict the applicability of these methods. As a motivational example, let us define function f on $D = [-\frac{1}{2}, \frac{5}{2}]$ by

$$f(x) = \begin{cases} x^3 \ln x^2 + x^5 - x^4, & x \neq 0 \\ 0, & x = 0 \end{cases}$$

Choose $x^* = 1$. We have that

$$\begin{aligned} f'(x) &= 3x^2 \ln x^2 + 5x^4 - 4x^3 + 2x^2, & f'(1) &= 3, \\ f''(x) &= 6x \ln x^2 + 20x^3 - 12x^2 + 10x \\ f'''(x) &= 6 \ln x^2 + 60x^2 - 24x + 22. \end{aligned}$$

Then, obviously, function f''' is unbounded on D . In the present paper we only use hypotheses on the first Fréchet derivative. This way we expand the applicability of method (2).

The rest of the paper is organized as follows: Section 2 contains the local convergence analysis of methods (2). The numerical examples are presented in the concluding Section 3.

2. LOCAL CONVERGENCE ANALYSIS

We present the local convergence analysis of method (2) in this section. Let $L_0 > 0, L > 0$ and $M \geq 1$. It is convenient for the local convergence analysis of method (2) to introduce some functions and parameters. Define functions g_1, g_2, h_2, g_3, h_3 on the interval $[0, \frac{1}{L_0})$ by

$$\begin{aligned} g_1(t) &= \frac{Lt}{2(1-L_0t)}, \\ g_2(t) &= \frac{1}{2(1-L_0t)} [Lt + 2Mg_1(t)], \\ h_2(t) &= g_2(t) - 1 \\ g_3(t) &= g_2(t) + \frac{M^3 g_1^2(t)(2+g_1(t))}{(1-\frac{L_0}{2}t)^2(1-L_0t)}, \\ h_3(t) &= g_3(t) - 1 \end{aligned}$$

and parameter

$$r_A = \frac{2}{2L_0 + L}.$$

We have that $h_2(0) = -1 < 0$ and $h_2(r_A) = \frac{M}{1 - L_0 r_A} > 0$, since $\frac{L r_A}{2(1 - L_0 r_A)} = 1$ and $1 - L_0 r_A > 0$. It then follows from the intermediate value theorem that function h_2 has zeros in the interval $(0, r_A)$. Denote by r_2 the smallest such zero. Similarly, we have that $h_3(0) = -1 < 0$ and $h_3(r_2) > 0$. Denote by r the smallest zero of function h_3 in the interval $(0, r_2)$. Then, for each $t \in [0, r)$ we have

$$(3) \quad 0 \leq g_1(t) < 1,$$

$$(4) \quad 0 \leq g_2(t) < 1,$$

and

$$(5) \quad 0 \leq g_3(t) < 1.$$

Denote by $U(v, \rho), \bar{U}(v, \rho)$ the open and closed balls in S , respectively, with center $v \in S$ and of radius $\rho > 0$. Next, we show the following local convergence result for method (2) using the preceding notation.

THEOREM 1. *Let $F : D \subseteq S \rightarrow S$ be a differentiable function. Suppose that there exist $x^* \in D$, $L_0 > 0, L > 0$ and $M \geq 1$ such that for each $x, y \in D$*

$$(6) \quad F(x^*) = 0, \quad F'(x^*) \neq 0,$$

$$(7) \quad |F'(x^*)^{-1}(F'(x) - F'(x^*))| \leq L_0|x - x^*|,$$

$$(8) \quad |F'(x^*)^{-1}(F'(x) - F'(y))| \leq L|x - y|,$$

$$(9) \quad |F'(x^*)^{-1}F'(x)| \leq M,$$

and

$$(10) \quad \bar{U}(x^*, (1 + M_0)r) \subseteq D,$$

where r is defined above [Theorem 1](#). Then, the sequence $\{x_n\}$ generated by method (2) for $x_0 \in U(x^*, r) - \{x^*\}$ is well defined, remains in $U(x^*, r)$ for each $n = 0, 1, 2, \dots$ and converges to x^* . Moreover, the following estimates hold

$$(11) \quad |y_n - x^*| \leq g_1(|x_n - x^*|)|x_n - x^*| < |x_n - x^*| < r,$$

$$(12) \quad |z_n - x^*| \leq g_2(|x_n - x^*|)|x_n - x^*| < |x_n - x^*|$$

and

$$(13) \quad |x_{n+1} - x^*| \leq g_3(|x_n - x^*|)|x_n - x^*| < |x_n - x^*|,$$

where the “ g ” functions are defined above in [Theorem 1](#). Furthermore, if that there exists $T \in [r, \frac{2}{L_0})$ such that $\bar{U}(x^*, T) \subset D$, then the limit point x^* is the only solution of equation $F(x) = 0$ in $\bar{U}(x^*, T)$.

Proof. We shall use induction to show estimates (11)–(13). Using the hypothesis $x_0 \in U(x^*, r) - \{x^*\}$, (7) and the definition of r , we have that

$$(14) \quad |F'(x^*)^{-1}(F'(x_0) - F'(x^*))| \leq L_0|x_0 - x^*| < L_0r < 1.$$

It follows from (14) and the Banach Lemma on invertible functions [3, 4, 19, 20, 22, 23] that $F'(x_0) \neq 0$ and

$$(15) \quad |F'(x_0)^{-1}F'(x^*)| \leq \frac{1}{1-L_0|x_0-x^*|} < \frac{1}{1-L_0r}.$$

Hence, y_0 and x_0 are well defined. We also get from (2), (3), (8) and (15) that

$$(16) \quad |y_0 - x^*| \leq |F'(x_0)^{-1}F'(x^*)| \int_0^1 F'(x^*)^{-1}(F'(x^* + \theta(x_0 - x^*)) - F'(x_0))(x_0 - x^*) d\theta \\ \leq \frac{L|x_0 - x^*|^2}{2(1-L_0|x_0 - x^*|)}$$

$$(17) \quad = g_1(|x_0 - x^*|)|x_0 - x^*| < |x_0 - x^*| < r,$$

which shows (11) for $n = 0$ and $y_0 \in U(x^*, r)$. We can write

$$(18) \quad F(x_0) - F(x^*) = \int_0^1 F'(x^* + \theta(x_0 - x^*))(x_0 - x^*) d\theta.$$

Then, by (9), (18) we obtain that

$$(19) \quad |F'(x^*)^{-1}F(x_0)| \leq \left| \int_0^1 F'(x^*)^{-1}F'(x^* + \theta(x_0 - x^*))(x_0 - x^*) d\theta \right| \\ \leq M|x_0 - x^*|,$$

where we also used $|x^* + \theta(x_0 - x^*) - x^*| = \theta|x_0 - x^*| < r$. That is $x^* + \theta(x_0 - x^*) - x^* \in U(x^*, r)$. We also have that

$$(20) \quad |F'(x^*)^{-1}F(y_0)| \leq M|y_0 - x^*| \leq Mg_1(|x_0 - x^*|)|x_0 - x^*|.$$

Then, by the second substep of method (2) for $n = 0$, (4), (6) and (20) we get that

$$(21) \quad |z_0 - x^*| \leq |x_0 - x^* - F'(x_0)^{-1}F(x_0)| \\ + |F'(x_0)^{-1}F'(x^*)||F'(x^*)^{-1}F(y_0)| \\ \leq \frac{L|x_0 - x^*|^2}{2(1-L_0|x_0 - x^*|)} + \frac{M|y_0 - x^*|}{1-L_0|x_0 - x^*|} \\ \leq \frac{(L|x_0 - x^*| + 2Mg_1(|x_0 - x^*|))|x_0 - x^*|}{2(1-L_0|x_0 - x^*|)} \\ = g_2(|x_0 - x^*|)|x_0 - x^*| < |x_0 - x^*| < r,$$

which shows (12) for $n = 0$ and $z_0 \in U(x^*, r)$. Next, we show that $F(x_0) \neq 0$. Using (7), we get that

$$|(F'(x^*)(x_0 - x^*))^{-1}[F(x_0) - F(x^*) - F'(x^*)(x_0 - x^*)]| \\ \leq |x_0 - x^*|^{-1} \left| \int_0^1 F'(x^*)^{-1}(F'(x^* + \theta(x_0 - x^*)) - F'(x^*))(x_0 - x^*) d\theta \right| \\ \leq |x_0 - x^*|^{-1} \frac{L_0|x_0 - x^*|^2}{2}$$

$$(22) \leq \frac{L_0|x_0-x^*|}{2} < \frac{L_0r}{2} < 1.$$

it follows from (22) that $F(x_0) \neq 0$ and

$$(23) \quad |F(x_0)^{-1}F'(x^*)| \leq \frac{1}{|x_0-x^*|(1-\frac{L_0}{2}|x_0-x^*|)}.$$

Hence x_1 is well defined. Then, using the last substep of method (2) for $n = 0$, (5), (15), (19), (20), (21) and (23) we get in turn that

$$\begin{aligned} |x_1 - x^*| &\leq |z_0 - x^*| + |F'(x^*)^{-1}F(y_0)|^2 |F'(x^*)^{-1}F(x_0)|^{-2} \\ &\quad \times |F'(x^*)^{-1}F'(x_0)|^{-1} (2|F'(x^*)^{-1}F(x_0)| \\ &\quad + |F'(x^*)^{-1}F(y_0)|) \\ &\leq g_2(|x_0 - x^*|)|x_0 - x^*| \\ &\quad + \frac{M^3|y_0-x^*|^2(2|x_0-x^*|+|y_0-x^*|)}{|x_0-x^*|^2(1-\frac{L_0}{2}|x_0-x^*|)^2(1-L_0|x_0-x^*|)} \\ &\leq [g_2(|x_0 - x^*|) \\ &\quad + \frac{M^3g_1^2(|x_0-x^*|)(2+g_1(|x_0-x^*|))}{(1-\frac{L_0}{2}|x_0-x^*|)^2(1-L_0|x_0-x^*|)}] |x_0 - x^*| \\ &= g_3(|x_0 - x^*|)|x_0 - x^*| < |x_0 - x^*| < r, \end{aligned}$$

which shows (13) for $n = 0$ and $x_1 \in U(x^*, r)$. By simply replacing x_0, y_0, z_0, x_1 by x_k, y_k, z_k, x_{k+1} in the preceding estimates we arrive at estimates (11)–(13). Using the estimate $|x_{k+1} - x^*| < |x_k - x^*| < r$, we deduce that $x_{k+1} \in U(x^*, r)$ and $\lim_{k \rightarrow \infty} x_k = x^*$. To show the uniqueness part, let $Q = \int_0^1 F'(y^* + \theta(x^* - y^*))d\theta$ for some $y^* \in \bar{U}(x^*, T)$ with $F(y^*) = 0$. Using (7) we get that

$$(24) \quad \begin{aligned} |F'(x^*)^{-1}(Q - F'(x^*))| &\leq \int_0^1 L_0|y^* + \theta(x^* - y^*) - x^*|d\theta \\ &\leq \int_0^1 (1 - \theta)|x^* - y^*|d\theta \leq \frac{L_0}{2}R < 1. \end{aligned}$$

It follows from (24) and the Banach Lemma on invertible functions that Q is invertible. Finally, from the identity $0 = F(x^*) - F(y^*) = Q(x^* - y^*)$, we deduce that $x^* = y^*$. \square

REMARK 2. 1. In view of (7) and the estimate

$$\begin{aligned} \|F'(x^*)^{-1}F'(x)\| &= \|F'(x^*)^{-1}(F'(x) - F'(x^*)) + I\| \\ &\leq 1 + \|F'(x^*)^{-1}(F'(x) - F'(x^*))\| \leq 1 + L_0\|x - x^*\| \end{aligned}$$

condition (9) can be dropped and M can be replaced by

$$M(t) = 1 + L_0t.$$

2. The results obtained here can be used for operators F satisfying autonomous differential equations [3] of the form

$$F'(x) = P(F(x))$$

where P is a continuous operator. Then, since $F'(x^*) = P(F(x^*)) = P(0)$, we can apply the results without actually knowing x^* . For example, let $F(x) = e^x - 1$. Then, we can choose: $P(x) = x + 1$.

3. The radius r_A was shown by us to be the convergence radius of Newton's method [2]–[4]

$$(25) \quad x_{n+1} = x_n - F'(x_n)^{-1}F(x_n) \text{ for each } n = 0, 1, 2, \dots$$

under the conditions (7) and (8). It follows from the definition of r that the convergence radius r of the method (2) cannot be larger than the convergence radius r_A of the second order Newton's method (25). As already noted in [3, 4] r_A is at least as large as the convergence ball given by Rheinboldt [27]

$$(26) \quad r_R = \frac{2}{3L}.$$

In particular, for $L_0 < L$ we have that

$$r_R < r$$

and

$$\frac{r_R}{r_A} \rightarrow \frac{1}{3} \text{ as } \frac{L_0}{L} \rightarrow 0.$$

That is our convergence ball r_A is at most three times larger than Rheinboldt's. The same value for r_R was given by Traub [28].

4. It is worth noticing that method (2) is not changing when we use the conditions of Theorem 1 instead of the stronger conditions used in [1, 5, 12]–[28]. Moreover, we can compute the computational order of convergence (COC) defined by

$$\xi = \ln \left(\frac{\|x_{n+1} - x^*\|}{\|x_n - x^*\|} \right) / \ln \left(\frac{\|x_n - x^*\|}{\|x_{n-1} - x^*\|} \right)$$

or the approximate computational order of convergence

$$\xi_1 = \ln \left(\frac{\|x_{n+1} - x_n\|}{\|x_n - x_{n-1}\|} \right) / \ln \left(\frac{\|x_n - x_{n-1}\|}{\|x_{n-1} - x_{n-2}\|} \right).$$

This way we obtain in practice the order of convergence in a way that avoids the bounds involving estimates using estimates higher than the first Fréchet derivative of operator F .

3. NUMERICAL EXAMPLES

We present numerical examples in this section.

EXAMPLE 3. Let $D = [-\infty, +\infty]$. Define function f of D by

$$(27) \quad f(x) = \sin(x).$$












Then we have for $x^* = 0$ that $L_0 = L = M = 1, \alpha = 1$. The parameters are $r_A = 0.6667, r_2 = 0.6667, r = 0.3991$ and $\xi_1 = 5.1010$.









EXAMPLE 4. Let $D = [-1, 1]$. Define function f of D by

$$(28) \quad f(x) = e^x - 1.$$

Using (28) and $x^* = 0$, we get that $L_0 = e - 1 < L = M = e, \alpha = 1$. The parameters are $r_A = 0.3249, r_2 = 0.1458, r = 0.0699$ and $\xi_1 = 3.9088$. \square

REFERENCES

- [1] S. AMAT, M.A. HERNÁNDEZ, N. ROMERO, *A modified Chebyshev's iterative method with at least sixth order of convergence*, Appl. Math. Comput., **206** (2008), 164–174, <https://doi.org/10.1016/j.amc.2008.08.050>. 
- [2] S. AMAT, S. BUSQUIER, S. PLAZA, *Dynamics of the King's and Jarratt iterations*, Aeq. Math., **69** (2005), 212–213, <https://doi.org/10.1007/s00010-004-2733-y>. 
- [3] I.K. ARGYROS, *Convergence and Application of Newton-type Iterations*, Springer, 2008.
- [4] I.K. ARGYROS, SAID HILOUT, *Computational Methods in Nonlinear Analysis*, World Scientific Publ. Co., New Jersey, USA, 2013.
- [5] I.K. ARGYROS, D. CHEN, Q. QUIAN, *The Jarratt method in Banach space setting*, J. Comput. Appl. Math., **51** (1994), pp. 103–106, [https://doi.org/10.1016/0377-0427\(94\)90093-0](https://doi.org/10.1016/0377-0427(94)90093-0). 
- [6] V. CANDELA, A. MARQUINA, *Recurrence relations for rational cubic methods I: The Halley method*, Computing, **44** (1990), 169–184, <https://doi.org/10.1007/bf02241866>. 
- [7] J. CHEN, *Some new iterative methods with three-order convergence*, Appl. Math. Comput., **181** (2006), 1519–1522, <https://doi.org/10.1016/j.amc.2006.02.037>. 
- [8] C. CHUN, B. NETA, M. SCOTT, *Basins of attraction for optimal eighth order methods to find simple roots of nonlinear equations*, Appl. Math. Comput., **227** (2014), pp. 567–592, <https://doi.org/10.1016/j.amc.2013.11.017>. 
- [9] A. CORDERO, J. TORREGROSA, *Variants of Newton's method using fifth order quadrature formulas*, Appl. Math. Comput., **190** (2007), pp. 686–698, <https://doi.org/10.1016/j.amc.2007.01.062>. 
- [10] A. CORDERO, J. MAIMO, J. TORREGROSA, M.P. VASSILEVA, P. VINDEL, *Chaos in King's iterative family*, Appl. Math. Lett., **26** (2013), pp. 842–848, <https://doi.org/10.1016/j.aml.2013.03.012>. 
- [11] A. CORDERO, A. MAGRENAN, C. QUEMADA, J.R. TORREGROSA, *Stability study of eight-order iterative methods for solving nonlinear equations*, J. Comput. Appl. Math., (to appear).
- [12] A. CORDERO, J.L. HUESO, E. MARTINEZ, J.R. TORREGROSSA, *Steffensen type methods for solving non-linear equations*, J. Comput. Appl. Math., **236** (2012), pp. 3058–3064, <https://doi.org/10.1016/j.cam.2010.08.043>. 
- [13] A. CORDERO, J.L. HUESO, E. MARTINEZ, J.R. TORREGROSSA, *New modifications of Potra-Ptak's method with optimal fourth and eight orders of convergence*, J. Comput. Appl. Math., **234** (2010), pp. 2969–2976, <https://doi.org/10.1016/j.cam.2010.04.009>. 
- [14] J.A. EZQUERRO, M.A. HERNÁNDEZ, *A uniparametric Halley-type iteration with free second derivative*, Int. J. Pure Appl. Math., **6** (2003), pp. 99–110.
- [15] J.A. EZQUERRO, M.A. HERNÁNDEZ, *New iterations of R-order four with reduced computational cost*, BIT Numer. Math., **49** (2009), pp. 325–342, <https://doi.org/10.1007/s10543-009-0226-z>. 

- [16] M. FRONTINI, E. SORMANI, *Some variants of Newton's method with third order convergence*, Appl. Math. Comput., **140** (2003), pp. 419–426, [https://doi.org/10.1016/s0096-3003\(02\)00238-2](https://doi.org/10.1016/s0096-3003(02)00238-2). 
- [17] J.M. GUTIÉRREZ, M.A. HERNÁNDEZ, *Recurrence relations for the super-Halley method*, Comput. Math. Appl., **36** (1998) 7, pp. 1–8, [https://doi.org/10.1016/s0898-1221\(98\)00168-0](https://doi.org/10.1016/s0898-1221(98)00168-0). 
- [18] M.A. HERNÁNDEZ, M.A. SALANOVA, *Sufficient conditions for semilocal convergence of a fourth order multipoint iterative method for solving equations in Banach spaces*, Southwest J. Pure Appl. Math., **1** (1999), pp. 29–40.
- [19] J.P. JAISWAL, *A new third-order derivative free method for solving nonlinear equations*, Universal J. Appl. Math., **1** (2013), pp. 131–135, <https://doi.org/10.13189/ujam.2013.010215>. 
- [20] R.F. KING, *A family of fourth-order methods for nonlinear equations*, SIAM. J. Numer. Anal., **10** (1973), pp. 876–879, <https://doi.org/10.1137/0710072>. 
- [21] A.K. MAHESHWARI, *A fourth order iterative method for solving nonlinear equations*, Appl. Math. Comput., **211** (2009), pp. 383–391, <https://doi.org/10.1016/j.amc.2009.01.047>. 
- [22] S.K. PARHI, D.K. GUPTA, *Semi-local convergence of a Stirling-like method in Banach spaces*, Int. J. Comput. Methods, **7** (2010), pp. 215–228, <https://doi.org/10.1142/s0219876210002210>. 
- [23] M.S. PETKOVIC, B. NETA, L. PETKOVIC, J. DŽUNIČ, *Multipoint Methods for Solving Nonlinear Equations*, Elsevier, 2013.
- [24] F.A. POTRA, V. PTAK, *Nondiscrete Induction and Iterative Processes*, Research Notes in Mathematics, vol. 103, Pitman Publ., Boston, MA, 1984.
- [25] L.B. RALL, *Computational Solution of Nonlinear Operator Equations*, Robert E. Krieger, New York, 1979.
- [26] H. REN, Q. WU, W. BI, *New variants of Jarratt method with sixth-order convergence*, Numer. Algor., **52** (2009), pp. 585–603, <https://doi.org/10.1007/s11075-009-9302-3>. 
- [27] W.C. RHEINBOLDT, *An adaptive continuation process for solving systems of nonlinear equations*, in: Mathematical models and numerical methods (A.N.Tikhonov *et al.* eds.) pub.3, (19), pp. 129–142 Banach Center, Warsaw Poland.
- [28] J.F. TRAUB, *Iterative Methods for the Solution of Equations*, Prentice Hall, Englewood Cliffs, New Jersey, USA, 1964.
- [29] X. WANG, J. KOU, *Convergence for modified Halley-like methods with less computation of inversion*, J. Diff. Eq. Appl., **19** (2013) 9, pp. 1483–1500, <https://doi.org/10.1080/10236198.2012.761979>. 

Received by the editors: May 5, 2015; accepted: June 2, 2016; published online: November 8, 2021.