

ON AN APPROXIMATION OPERATOR
AND ITS LIPSCHITZ CONSTANT

MARIA CRĂCIUN*

Abstract. In this note we consider an approximation operator of Kantorovich type in which expression appears a basic sequence for a delta operator and a Sheffer sequence for the same delta operator. We give a convergence theorem for this operator and we find its Lipschitz constant.

MSC 2000. 41A36.

Keywords. approximation operators of Kantorovich type, Sheffer sequences, Lipschitz constants.

1. INTRODUCTION

In this section we will remind some basic notions and results.

Let P be the linear space of all polynomials with real coefficients.

A polynomial sequence is a sequence of polynomials (p_n) with $\deg p_n = n$ for all $n \in \mathbb{N}$.

A sequence of binomial type (a binomial sequence) is a polynomial sequence which satisfies the binomial identity

$$p_n(x+y) = \sum_{k=0}^n \binom{n}{k} p_k(x) p_{n-k}(y)$$

for all real x, y and $n = 0, 1, 2, \dots$.

The shift operator $E^a : P \rightarrow P$ is defined by $E^a p(x) = p(x+a)$.

A linear operator T with $TE^a = E^aT$ for all real a is called a shift invariant operator.

We recall that if T_1 and T_2 are shift invariant operators then $T_1T_2 = T_2T_1$.

A delta operator is a shift invariant operator for which $Qx = \text{const.} \neq 0$.

A polynomial sequence (p_n) is called a basic sequence for a delta operator Q if $p_0(x) = 1$, $p_n(0) = 0$ and $Qp_n = np_{n-1}$, $n = 1, 2, \dots$.

PROPOSITION 1. [9]. i) *Every delta operator has a unique basic sequence.*

ii) *A polynomial sequence is a binomial sequence if and only if it is a basic sequence for a delta operator Q .*

The Pincherle derivative of an operator T is defined by $T' = TX - XT$, where X is the multiplication operator, $Xp(x) = xp(x)$.

*"T. Popoviciu" Institute of Numerical Analysis, P.O. Box 68-1, 3400 Cluj-Napoca, Romania, e-mail: craciun@ictp.acad.ro.

The Pincherle derivative of a shift invariant operator is also a shift invariant operator and the Pincherle derivative of a delta operator is an invertible operator.

A polynomial sequence $(s_n)_{n \geq 0}$ is called a Sheffer sequence relative to a delta operator Q if $s_0(x) = \text{const} \neq 0$ and $Qs_n = ns_{n-1}$, $n = 1, 2, \dots$.

An Appell sequence is a Sheffer sequence relative to the derivative D .

PROPOSITION 2. [9]. *Let Q be a delta operator with the basic sequence (p_n) and (s_n) a polynomial sequence. The following statements are equivalent:*

- i) s_n is a Sheffer set relative to Q .
- ii) There exists an invertible shift invariant operator S such that $s_n(x) = S^{-1}p_n(x)$.
- iii) For all $x, y \in \mathbb{R}$ and $n = 0, 1, 2, \dots$ the following identity holds:

$$s_n(x+y) = \sum_{k=0}^n \binom{n}{k} p_k(x) s_{n-k}(y).$$

2. AN APPROXIMATION OPERATOR OF KANTOROVICH TYPE

In our paper [3] we considered some linear approximation operators defined for all $f \in C[0, 1]$ and $x \in [0, 1]$ by

$$(1) \quad (L_n^{Q,S} f)(x) = \frac{1}{s_n(1)} \sum_{k=0}^n \binom{n}{k} p_k(x) s_{n-k}(1-x) f\left(\frac{k}{n}\right),$$

where (p_n) is the basic sequence for a delta operator Q and (s_n) is a Sheffer sequence for the same delta operator, $s_n(1) \neq 0$, $\forall n \in \mathbb{N}$, $s_n = S^{-1}p_n$ with S an invertible shift invariant operator.

We remind that if $p'_k(0) \geq 0$ and $s_k(0) \geq 0$ for $n = 0, 1, 2, \dots$ then the operator $L_n^{Q,S}$ defined by (1) is positive.

In this note we want to introduce an integral operator of Kantorovich type of the form

$$(2) \quad (K_n^{Q,S} f)(x) = \frac{(n+1)}{s_n(1)} \sum_{k=0}^n \binom{n}{k} p_k(x) s_{n-k}(1-x) \int_{\frac{k}{n+1}}^{\frac{k+1}{n+1}} f(t) dt,$$

where $f \in L_1([0, 1])$, $x \in [0, 1]$.

We mention that for $S = I$ (that means $s_n = p_n$) these operators were considered by O. Agratini in [1] and V. Miheşan in [6].

We recall that the expressions of the operator $L_n^{Q,S}$ on the test functions $e_k(x) = x^k$, $k = 0, 2$ are (see [3]):

$$\begin{aligned} (L_n^{Q,S} e_0)(x) &= e_0(x), \\ (L_n^{Q,S} e_1)(x) &= a_n e_1(x), \\ (L_n^{Q,S} e_2)(x) &= b_n x^2 + x(a_n - b_n - c_n), \end{aligned}$$

where

$$a_n = \frac{((Q')^{-1}s_{n-1})(1)}{s_n(1)}, \quad b_n = \frac{n-1}{n} \frac{((Q')^{-2}s_{n-2})(1)}{s_n(1)}, \quad c_n = \frac{n-1}{n} \frac{((Q')^{-2}(S^{-1})'Ss_{n-2})(1)}{s_n(1)}$$

and Q' is the Pincherle derivative of Q .

LEMMA 3. *If $K_n^{Q,S}$ is the linear operator defined by (2) then:*

$$\begin{aligned} (K_n^{Q,S}e_0)(x) &= e_0(x), \\ (K_n^{Q,S}e_1)(x) &= \frac{n}{n+1}a_n e_1(x) + \frac{1}{2(n+1)}, \\ (K_n^{Q,S}e_2)(x) &= \frac{1}{(n+1)^2} \left\{ x^2 n^2 b_n + x[n^2(a_n - b_n - c_n) + na_n] + \frac{1}{3} \right\}. \end{aligned}$$

Proof. If we denote $s_{n,k}(x) = \frac{1}{s_n(1)} \binom{n}{k} p_k(x) s_{n-k}(1-x)$ we have

$$\begin{aligned} (K_n^{Q,S}e_0)(x) &= (n+1) \sum_{k=0}^n s_{n,k}(x) \left(\frac{k+1}{n+1} - \frac{k}{n+1} \right) = 1 = e_0(x), \\ (K_n^{Q,S}e_1)(x) &= \frac{n}{n+1} \sum_{k=0}^n s_{n,k}(x) \frac{k}{n} + \frac{1}{2(n+1)} \sum_{k=0}^n s_{n,k}(x) \\ &= \frac{n}{n+1} (L_n^{Q,S}e_1)(x) + \frac{1}{2(n+1)} (L_n^{Q,S}e_0)(x) \\ &= \frac{n}{n+1} a_n e_1(x) + \frac{1}{2(n+1)}, \\ (K_n^{Q,S}e_2)(x) &= \frac{n^2}{(n+1)^2} (L_n^{Q,S}e_2)(x) + \frac{n}{(n+1)^2} (L_n^{Q,S}e_1)(x) \\ &\quad + \frac{1}{3(n+1)^2} (L_n^{Q,S}e_0)(x) \\ &= \frac{1}{(n+1)^2} \left\{ x^2 n^2 b_n + x[n^2(a_n - b_n - c_n) + na_n] + \frac{1}{3} \right\}. \quad \square \end{aligned}$$

From this Lemma, the central moments of $K_n^{Q,S}$ defined by $\Omega_{n,k}(x) = K_n^{Q,S}((e_1 - x e_0)^k, x)$, $k \in \mathbb{N}$ are

$$\begin{aligned} \Omega_{n,0}(x) &= 1, \\ \Omega_{n,1}(x) &= \frac{1}{n+1} \left[x(na_n - (n+1)) + \frac{1}{2} \right] \quad \text{and} \\ \Omega_{n,2}(x) &= \frac{1}{(n+1)^2} \left\{ x^2 [n^2(b_n - 2a_n + 1) + 2n(1 - a_n)] \right. \\ &\quad \left. + x [n^2(a_n - b_n - c_n) + n(a_n - 1) - 1] + \frac{1}{3} \right\}. \end{aligned}$$

THEOREM 4. *Let $K_n^{Q,S}$ be the linear operator defined by (2) with $p'_k(0) \geq 0$ and $s_k(0) \geq 0, \forall k \in \mathbb{N}$.*

- i) *If $f \in C[0,1]$, $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n = 1$ then $K_n^{Q,S}$ converges uniformly to f .*
- ii) *If $f \in L_p[0,1]$, $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n = 1$ then $\|K_n^{Q,S}f - f\|_p = 0$.*

Proof. In [3] we proved that if $p'_k(0) \geq 0$ and $s_k(0) \geq 0, \forall k \in \mathbb{N}$ then $0 \leq c_n \leq \min\{(1-b_n)/2, a_n - a_n^2\}$, so from $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n = 1$ we have $\lim_{n \rightarrow \infty} c_n = 0$. Using Lemma 3 it results that $\lim_{n \rightarrow \infty} (K_n^{Q,S} e_i)(x) = e_i(x)$ for $i = 0, 1, 2$, and applying the convergence criterion of Bohman-Korovkin we obtain the first affirmation.

The second assertion follows immediately because the Korovkin subspaces in $C[0, 1]$ are also Korovkin subspaces in $L_p[0, 1]$. \square

3. LIPSCHITZ CONSTANTS FOR $L_n^{Q,S}$ AND $K_n^{Q,S}$

In this section we want to find the Lipschitz constants for $L_n^{Q,S}$ and $K_n^{Q,S}$ if $f \in Lip_M \alpha$.

In [2] B.M. Brown, D. Elliot and D.F. Paget proved that the Bernstein operator ($B_n = L_n^{D,I}$) preserves the Lipschitz constant of the function f for $\alpha \in (0, 1]$. V. Miheşan showed in [6] that all positive binomial operators (which can be obtained by $L_n^{Q,S}$ when $S = I$) preserve the Lipschitz constant of the function f for $\alpha \in (0, 1]$ and if $f \in Lip_M^*(\alpha, [0, 1])$ then $L_n^{Q,I} f \in Lip_{2M}^*(\alpha, [0, 1])$, where

$$Lip_M^*(\alpha, [0, 1]) = \{f \in C[0, 1], \omega_2(f, h) \leq Mh^\alpha, 0 < h \leq \frac{1}{2}\}.$$

THEOREM 5. *If $f \in Lip_M \alpha, \alpha \in (0, 1]$, then $L_n^{Q,S} f \in Lip_{M\alpha_n^\alpha}$.*

Proof. Let $x \leq y$ be any two points of $[0, 1]$. Using the binomial identity for p_n we can write

$$\begin{aligned} (L_n^{Q,S} f)(y) &= \frac{1}{s_n(1)} \sum_{j=0}^n \binom{n}{j} p_j(x + (y-x)) s_{n-j}(1-y) f\left(\frac{j}{n}\right) \\ &= \frac{1}{s_n(1)} \sum_{j=0}^n \binom{n}{j} s_{n-j}(1-y) f\left(\frac{j}{n}\right) \sum_{k=0}^j \binom{j}{k} p_k(x) p_{j-k}(y-x). \end{aligned}$$

If we change the order of summation and note $j - k = l$ then we obtain

$$\begin{aligned} (3) \quad (L_n^{Q,S} f)(y) &= \\ &= \frac{1}{s_n(1)} \sum_{k=0}^n \sum_{l=0}^{n-k} \frac{n!}{k!l!(n-k-l)!} p_k(x) p_l(y-x) s_{n-k-l}(1-y) f\left(\frac{k+l}{n}\right), \\ (L_n^{Q,S} f)(x) &= \frac{1}{s_n(1)} \sum_{k=0}^n \binom{n}{k} p_k(x) s_{n-k}((y-x) + (1-y)) f\left(\frac{k}{n}\right) \\ &= \frac{1}{s_n(1)} \sum_{k=0}^n \binom{n}{k} p_k(x) f\left(\frac{k}{n}\right) \sum_{l=0}^{n-k} \binom{n-k}{l} p_l(y-x) s_{n-k-l}(1-y) f\left(\frac{k}{n}\right), \\ (4) \quad (L_n^{Q,S} f)(x) &= \\ &= \frac{1}{s_n(1)} \sum_{k=0}^n \sum_{l=0}^{n-k} \frac{n!}{k!l!(n-k-l)!} p_k(x) p_l(y-x) s_{n-k-l}(1-y) f\left(\frac{k}{n}\right). \end{aligned}$$

From (3) and (4) we have

$$\begin{aligned} & \left| \left(L_n^{Q,S} f \right) (y) - \left(L_n^{Q,S} f \right) (x) \right| = \\ & = \frac{1}{s_n(1)} \sum_{k=0}^n \sum_{l=0}^{n-k} \frac{n!}{k!l!(n-k-l)!} p_k(x) p_l(y-x) s_{n-k-l}(1-y) \left| f\left(\frac{k+l}{n}\right) - f\left(\frac{k}{n}\right) \right|. \end{aligned}$$

Because $f \in Lip_M \alpha$ we have $\left| f\left(\frac{k+l}{n}\right) - f\left(\frac{k}{n}\right) \right| \leq M \left(\frac{l}{n}\right)^\alpha$ so we obtain

$$\begin{aligned} & \left| \left(L_n^{Q,S} f \right) (y) - \left(L_n^{Q,S} f \right) (x) \right| \leq \\ & \leq \frac{M}{s_n(1)} \sum_{k=0}^n \sum_{l=0}^{n-k} \frac{n!}{k!l!(n-k-l)!} p_k(x) p_l(y-x) s_{n-k-l}(1-y) \left(\frac{l}{n}\right)^\alpha \\ & = \frac{M}{s_n(1)} \sum_{l=0}^n \sum_{k=0}^{n-l} \binom{n-l}{k} p_k(x) s_{n-k-l}(1-y) \binom{n}{l} p_l(y-x) \left(\frac{l}{n}\right)^\alpha \\ & = \frac{M}{s_n(1)} \sum_{l=0}^n \binom{n}{l} p_l(y-x) s_{n-l}(x+1-y) \left(\frac{l}{n}\right)^\alpha \\ & = M L_n^{Q,S}(x^\alpha; y-x). \end{aligned}$$

We remind that for a convex function f we have $f(a_n x) \leq (L_n^{Q,S} f)(x)$ (see [3]). Since the function $g(x) = -x^\alpha$, $\alpha \in (0, 1]$, is convex on $[0, 1]$ we obtain

$$\left(L_n^{Q,S} x^\alpha; y-x \right) \leq (a_n(y-x))^\alpha$$

and we get

$$\left| \left(L_n^{Q,S} f \right) (y) - \left(L_n^{Q,S} f \right) (x) \right| \leq M a_n^\alpha (y-x)^\alpha.$$

Therefore $L_n^{Q,S} f \in Lip_{M a_n^\alpha} \alpha$. \square

THEOREM 6. *If $f \in Lip_M \alpha$, $\alpha \in (0, 1]$, then $K_n^{Q,S} f \in Lip_{N_n} \alpha$, where $N_n = M \left(\frac{n a_n}{n+1}\right)^\alpha$.*

Proof. We can write $K_n^{Q,S} f = L_n^{Q,S} h_n$, where

$$\begin{aligned} h_n(x) &= \int_0^1 f\left(\frac{t+nx}{n+1}\right) dt, \\ |h_n(x) - h_n(y)| &= \left| \int_0^1 \left[f\left(\frac{t+nx}{n+1}\right) - f\left(\frac{t+ny}{n+1}\right) \right] dt \right| \\ &\leq M \left| \frac{t+nx}{n+1} - \frac{t+ny}{n+1} \right|^\alpha \leq M \left(\frac{n}{n+1}\right)^\alpha |x-y|^\alpha. \end{aligned}$$

So, $f \in Lip_M \alpha$ implies $h_n \in Lip_{M \left(\frac{n}{n+1}\right)^\alpha} \alpha$. From $K_n^{Q,S} f = L_n^{Q,S} h_n$ and the previous theorem we obtain the conclusion. \square

REFERENCES

- [1] AGRATINI, O., *On a certain class of approximation operators*, Pure Math. Appl., **11**, pp. 119–127, 2000.
- [2] BROWN, B. M., ELLIOT, D. and PAGET, D. F., *Lipschitz constants for the Bernstein polynomials of a Lipschitz continuous function*, J. Approx. Theory, **49**, pp. 196–199, 1987.
- [3] CRĂCIUN, M., *Approximation operators constructed by means of Sheffer sequences*, Rev. Anal. Numér. Théor. Approx., **30**, 2001, pp. 135–150. [✉](#)
- [4] LUPAŞ, L. and LUPAŞ, A., *Polynomials of binomial type and approximation operators*, Studia Univ. Babeş-Bolyai, Mathematica, **32**, pp. 61–69, 1987.
- [5] MANOLE, C., *Approximation operators of binomial type*, Univ. Cluj-Napoca, Research Seminar on Numerical and Statistical Calculus, Preprint no. 9, pp. 93–98, 1987.
- [6] MIHEŞAN, V., *Approximation of continuous functions by means of linear positive operators*, Ph.D. Thesis, Cluj-Napoca, 1997 (in Romanian).
- [7] MOLDOVAN, G., *Discrete convolutions and linear positive operators*, Ann. Univ. Sci. Budapest R. Eötvös, **15**, pp. 31–44, 1972.
- [8] POPOVICIU, T., *Remarques sur les polynomes binomiaux*, Bull. Soc. Math. Cluj, **6**, pp. 146–148, 1931.
- [9] ROTA, G.-C., KAHANER, D. and ODLYZKO, A., *On the foundations of combinatorial theory. VIII. Finite operator calculus*, J. Math. Anal. Appl., **42**, pp. 684–760, 1973.
- [10] SABLONNIÈRE, P., *Positive Bernstein-Sheffer operators*, J. Approx. Theory, **83**, pp. 330–341, 1995.
- [11] STANCU, D. D., *Approximation of functions by a new class of linear positive operators*, Rev. Roum. Math. Pures Appl., **13**, pp. 1173–1194, 1968.
- [12] STANCU, D. D., *On the approximation of functions by means of the operators of binomial type of Tiberiu Popoviciu*, Rev. Anal. Numér. Théor. Approx., **30**, pp. 95–105, 2001. [✉](#)
- [13] STANCU, D. D. and OCCORSIO, M. R., *On approximation by binomial operators of Tiberiu Popoviciu type*, Rev. Anal. Numér. Théor. Approx., **27**, pp. 167–181, 1998. [✉](#)

Received by the editors: September 27, 2001.