



ELSEVIER

Available online at www.sciencedirect.com



Computers and Mathematics with Applications 55 (2008) 1178–1180

An International Journal
**computers &
mathematics**
with applications

www.elsevier.com/locate/camwa

On the iterates of a class of summation-type linear positive operators

Octavian Agratini

Babeş-Bolyai University, Faculty of Mathematics and Computer Science, 400084 Cluj-Napoca, Romania

Received 6 February 2007; received in revised form 11 April 2007; accepted 18 April 2007

Abstract

This note is focused upon positive linear operators which preserve the quadratic test function. By using contraction principle, we study the convergence of their iterates.

© 2007 Elsevier Ltd. All rights reserved.

Keywords: Linear positive operator; Popoviciu–Bohman–Korovkin theorem; Contraction principle; Modulus of continuity; Iterates

1. Introduction

Many of the linear methods of approximation are given by a sequence of linear positive operators (*LPOs*) designed as follows

$$(A_n f)(x) = \sum_{k=0}^n a_{n,k}(x) f(x_{n,k}), \quad f \in C([a, b]), \quad x \in [c, d], \quad (1)$$

where every function $a_{n,k} \in C([c, d])$ is non-negative, $a = x_{n,0} < x_{n,1} < \dots < x_{n,n} = b$ forms a mesh of nodes and $a \leq c < d \leq b$. As usual, we consider that the Banach space $C(K)$, $K \subset \mathbb{R}$ compact interval, is endowed with the norm $\|\cdot\|_K$ of the uniform convergence. In accordance with Popoviciu–Bohman–Korovkin theorem, if $(\|e_i - A_n e_i\|_{[c,d]})_n$ tends to zero for $i \in \{0, 1, 2\}$, then $(A_n f)_n$ converges uniformly to f for each $f \in C([a, b])$. Here e_i stands for the monomial of i -degree.

King [1] constructed operators of Bernstein-type which reproduce the test functions e_0 and e_2 . Starting from a similar class of operators having the degree of exactness zero, our aim is to study the convergence of the iterates and some approximation properties of our class as well.

2. The construction

As regards the operators defined by (1), we assume that the following identities

$$\sum_{k=0}^n a_{n,k}(x) = 1, \quad \sum_{k=0}^n a_{n,k}(x) x_{n,k}^2 = x^2, \quad x \in [c, d], \quad n \in \mathbb{N}, \quad (2)$$

are fulfilled.

E-mail address: agratini@math.ubbcluj.ro.

A generalization to the m -dimensional case will be read as follows. Let K_m be a compact and convex subset of the space \mathbb{R}^m . Volkov [2] proved that the functions: $\mathbf{1}, pr_1, \dots, pr_m, \sum_{j=1}^m pr_j^2$, are test functions for $C(K_m)$. Here $pr_j, j = \overline{1, m}$, represent the canonical projections on K_m . Let $A_n : C(K_m) \rightarrow C(K_m)$ be such that

$$A_n \mathbf{1} = \mathbf{1} \quad \text{and} \quad A_n \left(\sum_{j=1}^m pr_j^2 \right) = \sum_{j=1}^m pr_j^2. \tag{3}$$

3. Results

By using the contraction principle we study the convergence of the iterates of the uni-dimensional operators A_n . We put $A_n^{m+1} = A_n \circ A_n^m, m \in \mathbb{N}$, and A_n^0 represents the identity operator of the space $C([a, b])$.

Theorem 3.1. *Let $A_n, n \in \mathbb{N}$, be defined by (1) and (2) such that $a = c < d = b, b \neq -a$ and $a_{n,0}(a) = a_{n,n}(b) = 1$. Set $u_n = \min_{x \in [a,b]} (a_{n,0}(x) + a_{n,n}(x))$. If $u_n > 0$, then the iterates sequence $(A_n^m)_{m \geq 1}$ verifies*

$$\lim_{m \rightarrow \infty} (A_n^m f)(x) = \frac{1}{b^2 - a^2} \left(f(a)b^2 - f(b)a^2 + (f(b) - f(a))x^2 \right), \tag{4}$$

uniformly with respect to x on $[a, b]$.

Considering the m -dimensional case of A_n described by (3), one has

$$\|A_n f - f\|_{K_m} \leq 2\omega(f; \sqrt{\mu_n}). \tag{5}$$

Here $\omega(f; \cdot)$ represents the modulus of continuity for the function f and

$$\mu_n := 2\|x\|^2 - 2 \sum_{i=1}^m x_i A_n(pr_i; x).$$

Proof. We define $X_{A,B} := \{f \in C([a, b]) \mid f(a) = A, f(b) = B\}, A \in \mathbb{R}, B \in \mathbb{R}$. Every $X_{A,B}$ is a closed subset of $C([a, b])$ and the system $X_{A,B}, (A, B) \in \mathbb{R} \times \mathbb{R}$, makes up a partition of this space. Since $a_{n,0}(a) = 1$, the first identity of (2) implies $a_{n,k}(a) = 0, k = 1, n$, consequently $(A_n f)(a) = f(a) = A$. Analogously, $(A_n f)(b) = f(b) = B$. These relations ensure that $X_{A,B}$ is an invariant subset of A_n for any $A \in \mathbb{R}$ and $B \in \mathbb{R}$.

Further on, we prove that the restriction of A_n at $X_{A,B}$ is a contraction for any $A \in \mathbb{R}$ and $B \in \mathbb{R}$. Indeed, if f and g belong to $X_{A,B}$ then, for each $a \leq x \leq b$, we can write

$$\begin{aligned} |(A_n f)(x) - (A_n g)(x)| &= \left| \sum_{k=1}^{n-1} a_{n,k}(x)(f - g)(x_{n,k}) \right| \leq \sum_{k=1}^{n-1} a_{n,k}(x) \|f - g\|_{[a,b]} \\ &= (1 - a_{n,0}(x) - a_{n,n}(x)) \|f - g\|_{[a,b]} \leq (1 - u_n) \|f - g\|_{[a,b]}, \end{aligned}$$

and consequently, $\|A_n f - A_n g\|_{[a,b]} \leq (1 - u_n) \|f - g\|_{[a,b]}$. On the other hand, the function

$$p_{A,B}^* = \frac{Ab^2 - Ba^2}{b^2 - a^2} e_0 + \frac{B - A}{b^2 - a^2} e_2$$

belongs to $X_{A,B}$. Since $A_n e_0 = e_0, A_n e_2 = e_2, p_{A,B}^*$ is a fixed point of A_n . For any $f \in C([a, b])$ one has $f \in X_{f(a),f(b)}$ and, by using the contraction principle, we get (4).

For the m -dimensional case, we can write

$$A_n(\|\cdot - x\|^2; x) = A_n \left(\sum_{i=1}^m (\cdot - x_i)^2; x \right) = 2\|x\|^2 - 2 \sum_{i=1}^m x_i A_n(pr_i; x),$$

where $\|\cdot\|$ stands for the Euclidean norm in \mathbb{R}^m . Taking into account a result established by Censor [3], Eq. (5), our relation (5) follows. \square

References

- [1] J.P. King, Positive linear operators which preserve x^2 , *Acta Math. Hungar.* 99 (3) (2003) 203–208.
- [2] V.I. Volkov, On the convergence of sequences of linear positive operators in the space of continuous functions of two variables, *Dokl. Akad. Nauk SSSR (N.S.)* 115 (1957) 17–19 (in Russian).
- [3] E. Censor, Quantitative results for positive linear approximation operators, *J. Approx. Theory* 4 (1971) 442–450.